

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

B5

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/395, 48/00, C07K 16/30, 16/46, C12N 5/20	A1	(11) International Publication Number: WO 96/37224 (43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: PCT/US96/07321 (22) International Filing Date: 21 May 1996 (21.05.96) (30) Priority Data: 08/449,911 25 May 1995 (25.05.95) US (71) Applicant: LUDWIG INSTITUTE FOR CANCER RE- SEARCH [CH/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). (72) Inventors: WELT, Sydney; 1345 Avenue of the Americas, New York, NY 10105 (US). OLD, Lloyd, J.; 1345 Avenue of the Americas, New York, NY 10105 (US). BARENDSSWAARD, Elsje; 1345 Avenue of the Americas, New York, NY 10105 (US). MONTALTO, Nicholas, J.; 1345 Avenue of the Americas, New York, NY 10105 (US). GURE, Ali, Osmay; 1345 Avenue of the Americas, New York, NY 10105 (US). (74) Agent: PASQUALINI, Patricia, A.; Felfe & Lynch, 805 Third Avenue, New York, NY 10022 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHODS OF TREATING COLON CANCER UTILIZING TUMOR-SPECIFIC ANTIBODIES (57) Abstract This invention relates to methods of reducing the effects of colon cancer tumors. Various agents are conjugated to monoclonal antibodies which are specific for colon cancer cells. The conjugates are administered to patients having colon cancer such that the effects of the cancer are reduced.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHODS OF TREATING COLON CANCER
UTILIZING TUMOR-SPECIFIC ANTIBODIES

Cross-Reference to Related Applications

5 This Application is a Continuation-In-Part of U.S. Patent
Application Serial No. 08/020,223 filed February 16, 1993,
pending; which is a Continuation Application of U.S. Patent
Application Serial No. 07/673,153 filed March 18, 1991,
abandoned; which is a Continuation Application of U.S. Patent
10 Application Serial No. 07/327,765 filed March 23, 1989,
abandoned; which is a Continuation-In-Part Application of U.S.
Patent Application Serial No. 07/118,411 filed November 6,
1987, abandoned; which is a Continuation Application of U.S.
Patent Application Serial No. 06/724,991 filed April 19, 1985,
15 abandoned.

FIELD OF THE INVENTION

 This invention relates to a method of reducing the
effects of colon cancer tumors, utilizing at least one
monoclonal antibody. Specifically, at least one monoclonal
20 antibody is utilized in conjunction with an anti-tumor drug,
a peptide which inhibits DNA tumor activity or a radioisotope
in the treatment of colorectal carcinoma. This invention
further relates to a method of delivering genetic material to
DNA of tumor cells and to a method of delivering anti-cancer
25 agents to nuclei of colon tumor cells, as well as to
monoclonal antibodies which are specific for A33 antigen; an
antigen found on colon cancer cells.

BACKGROUND OF THE INVENTION

 Colorectal carcinoma is a malignant neoplastic disease.
30 There is a high incidence of colorectal carcinoma in the
Western world, particularly in the United States. Tumors of
this type often metastasize through lymphatic and vascular
channels. Many patients with colorectal carcinoma eventually
die from this disease. In fact, it is estimated that 62,000
35 persons in the United States alone die of colorectal carcinoma
annually.

 To date, systemic therapies and chemotherapies have been

developed for the treatment of colorectal cancer. However, no therapies have exhibited sufficient anti-tumor activity to prolong the survival of colorectal carcinoma patients with metastatic disease with any degree of reliability. As a result, a need still exists to develop methods for the successful treatment of colorectal carcinoma.

It is therefore an object of this invention to provide methods for reducing the effects of colon cancers.

It is another object of this invention to provide a method of delivering genetic material to DNA of colon cancer cells.

It is a further object of this invention to provide a method of delivering anti-cancer agents to nuclei of colon cancer cells.

It is a still further object of this invention to provide antibodies, including monoclonal antibodies, humanized antibodies, chimeric antibodies, trimeric antibodies, heteromeric antibodies and single chain antibodies which are useful in the treatment of colon cancer.

SUMMARY OF THE INVENTION

This invention is directed to methods of reducing the effects of colon cancer comprising the administration of tumor-associated antibody conjugates. Specifically, at least one conjugate of an antibody which is associated with colon cancer cells and an anti-cancer drug is administered. Alternatively, at least one conjugate of an antibody which is specific for colon cancer cells and a peptide which inhibits DNA activity of said cells is administered in a pharmaceutically effective amount so as to reduce the effects of colon cancer. Another method of reducing the effects of colon cancer comprises administering a pharmaceutically effective amount of at least one conjugate of an antibody which is specific for said tumor and a radioisotope.

This invention is further directed to a method of delivering genetic material to colon cancer cells comprising contacting tumor cells with genetic material conjugated to an antibody which is internalized into the tumor cells, thereby

mediating integration of genetic material into DNA.

In addition, anti-cancer agents can be delivered to the nuclei (DNA) of colon cancer cells by contacting the cells with an anti-cancer agent conjugated to an antibody which is internalized into said cells.

This invention is also directed to antibodies, including monoclonal, humanized, chimeric, trimeric, heteromeric and single chain antibodies, which can be utilized to reduce the effects of colon cancers.

BRIEF DESCRIPTION OF THE DRAWINGS

The above brief description, as well as further objects and features of the present invention, will be more fully understood by reference to the following detailed description of the presently preferred, albeit illustrative, embodiments of the present invention when taken in conjunction with the accompanying drawings wherein:

Figure 1 is an autoradiograph which shows the accumulation of ^{131}I -mAb A33 in hepatic lesions but not in surrounding normal liver tissue;

Figure 2 is an autoradiograph which shows that radioisotope ^{131}I , when conjugated to mAb A33, is internalized into tumor cells and is not found in surrounding tissue;

Figure 3 represents the effect of ^{131}I -mAb A33 on carcinomatous lesions;

Figure 4 represents vectors for hA33 IgG1 and hA33 Fab' Δ cys expression;

Figure 5 represents heavy and light chain variable domain amino acid sequences for humanized A33 antibodies;

Figure 6 represents SDS polyacrylamide gel of humanized A33 IgG and TFM[™] under reducing and non-reducing conditions;

Figure 7 represents HPLC profiles of hA33 Fab', hA33 Fab' after cross-linking to TFM[™], and purified hA33 TFM[™];

Figure 8 represents a Scatchard plot of humanized and mouse A33 binding to SW1222 cells;

Figure 9 represents a competitive binding assay for hA33 Fab' Δ cys, IgG and TFM[™] binding to COLO 205 cells;

Figure 10 shows biodistribution of ^{90}Y -labelled humanized

A33 in nude mice bearing SW1222 tumor xenografts; and

Figure 11 shows pharmacokinetic profiles of ¹¹¹In hA33 IgG, TFM[™] and DFM[™] in cynomolgus monkeys.

DETAILED DESCRIPTION OF THE INVENTION

5 Murine monoclonal antibody A33, (referred to herein as mAb A33) is a monoclonal antibody of the IgG2a isotype which defines a cell-surface antigen referred to as antigen A33, which is present on greater than 95% of colorectal carcinomas and in normal intestinal mucosa (see Table 1). Among
10 colorectal carcinomas, A33 antigen is expressed with a homogeneous cell surface distribution in greater than 95% of antigen-positive lesions. Moreover, A33 antigen is found in colon cancers regardless of their degree of histologic differentiation and in primary as well as metastatic lesions,
15 including liver metastases.

TABLE 1

**Immunohistochemical Analysis of A33
Antigen Expression in Human Tumors**

	Tumor Type	A33-positive No. Cases	% positive cells
5	-----		
10	Carcinomas		
	Colorectal carcinoma		
	primary tumors	45/47	75-100 n = 40 25-50 n = 5
	metastases	23/25	75-100 n = 22 25-50 n = 1
15	Gastric carcinoma	14/30	
	signet ring type	12/12	75-100 n = 12
20	Esophageal carcinoma	2/8	
	intestinal type	1/1	
	mucinous type	1/1	
	Pancreatic carcinoma	0/8	
	Lung carcinoma	0/16	
	Breast carcinoma	0/19	
25	Renal carcinoma	0/16	
	Bladder carcinoma	0/19	
	Prostate carcinoma	0/4	
	Testicular carcinoma	0/4	
	Ovarian carcinoma	1/56	
30	Endometrial carcinoma	1/4	
	Thyroid carcinoma	0/4	
	Liver carcinoma	0/2	
	Larynx carcinoma	0/2	
	Mesothelioma	1/7	
35	Neuroendocrine carcinomas	0/8	
	Neuroectodermal tumors		
	Melanoma	1/10	
40	Gliomas	0/8	
	Neuroblastomas, Glqnb	0/10	
	Sarcomas		
	Leiomyosarcoma	0/7	
45	MFII	0/5	
	Fibrosarcoma	0/3	
	Liposarcoma	0/6	
	MPNT	0/6	
	Chondrosarcoma	0/14	
50	Others	0/9	
	Lymphomas	0/12	

Antigen A33 has not been detected in the sera of patients with antigen A33-positive colon cancers, in the tumoral secretions of mucinous colon cancers, nor in the supernatant of antigen A33-positive colon cancer cell lines. This suggests that antigen A33 is not a secreted molecule. In the normal gastrointestinal tract, antigen A33 is found in normal small and large intestinal mucosa, showing a uniform cell surface distribution throughout the crypts. A detailed survey of antigen A33 expression in greater than 300 tumors of diverse histologic types and in normal human adult and fetal tissues has shown the restricted distribution pattern of this antigen. As discussed hereinbelow, the inventors have found that among neoplastic tissues, antigen A33 is present in colon cancers and in a subset of gastric cancers showing intestinal metaplasia, but not in any of the other epithelial cancers tested. Antigen A33 was not found to be present in sarcomas, neuroectodermal tumors, or lymphoid neoplasms. These tissues were consistently found to be antigen A33-negative.

As disclosed in U.S. Patent No. 5,160,723, the entirety of which is incorporated herein by reference, monoclonal antibody A33 is specific for antigen A33 found on colon cancer cell surfaces, and is therefore specific for colon cancer. As a result, monoclonal antibody A33, and other antibodies which are antigen A33-specific, can be used for colon cancer diagnosis. The antibodies can be conjugated with anti-cancer agents, peptides or radioisotopes and utilized for colon tumor treatment.

Example I - Internalization of ^{125}I -mAb A33
in Heterotransplanted Human Colon Tumors in Mice

The A33 antigen has several features which make it useful for immunotherapy studies. It is present in more than 95% of colorectal cancers, including primary and metastatic lesions. It is homogeneously expressed in tumor nodules, and it is not secreted into the blood stream.

Further, clinical evaluation of ^{131}I - and ^{125}I -mAb A33 have shown exceptionally good targeting to metastatic colon cancer

sites and evidence for clinical responses with only limited toxicity. Biopsy-based quantitative dosimetry data for mAb A33 show that mAb A33 can be used for effective and safe radioimmunotherapy.

5 Monoclonal antibody A33 was radiolabelled with ^{125}I by procedures described in U.S. Patent No. 5,160,723 (the '723 Patent). The ^{125}I -labelled A33 monoclonal antibody is referred to herein as ^{125}I -mAb A33. ^{125}I -mAb A33 was administered to
10 nu/nu mice which were heterotransplanted with human colon tumors. The mice were subsequently studied for ^{125}I -mAb A33 localization. Parameters of radiolabelled monoclonal antibody A33 localization were evaluated in nu/nu mice using the A33-positive colon carcinoma cell line SW1222. This model is of special interest because much is known about the kinetics and
15 trafficking of mAb A33 in cultured SW1222 cells *in vitro*. Specifically, it is known that mAb A33 is rapidly internalized into SW1222 cells (approximately 33% in less than 1 hour); that the pathway of mAb A33 internalization is via macropinosomes generated from cell membrane ruffling; that on
20 average, SW1222 cells bind 800,000 molecules of mAb A33 per cell; that the $t_{1/2}$ of ^{125}I -mAb A33 retention by these cells is 12-14 hours; and that most of the internalized mAb A33 is released in intact immunoreactive form. The released mAb A33 is capable of rebinding to tumor cells, resulting in higher
25 uptake/retention of this monoclonal antibody. This occurs in both mice and humans (human data described hereinbelow).

The peak tumor uptake for ^{125}I -mAb A33 at 24-48 hours after injection was 35-40% injected dose/g. Tumor:liver ratios ranged from 30:1 to 40:1, and tumor:blood ratios ranged
30 from 10:1 to 15:1. These values showed some minor variation depending on the dose of monoclonal antibody administered, and localization parameters were found to be antigen-specific as tumor uptake could be blocked by the intravenous
administration of excess unlabelled mAb A33, but not with
35 unrelated negative control mAb. ^{125}I -mAb A33 was cleared from tumor xenografts with a $t_{1/2}$ of 60 hours. In contrast, mAb A33 radiolabelled with ^{111}In showed localization of 45%

5 injected dose/g tumor, and retention of the isotope was significantly longer (greater than 7 days). This was most likely due to cytoplasmic trapping of ^{111}In . This *in vivo* model of mAb A33 localization to tumor is of great value in evaluating the relative effectiveness of various antibody conjugates, and for dosimetry calculations involving a variety of isotopes.

10 Example II - Internalization
of ^{125}I -mAb A33 in Humans with Colon Cancer

15 The A33 antigenic system can be used for immunotherapy. It is particularly suitable for this use since there is rapid internalization of antigen A33-mAb A33 complexes into colon cancer cells. Further, mAb A33 conjugates, which have intracellular sites of action, have the ability to kill colon cancer cells *in vitro* and *in vivo*.

20 The inventors studied the localization and cytotoxic effects of ^{125}I -labelled mAb A33 in humans. ^{125}I radionuclide exerts its cytotoxic effects primarily through short-range Auger electrons, which are most effective when generated in close proximity ($<1\text{-}4\ \mu\text{m}$) to the cell nucleus. Accordingly, one of the expected benefits of ^{125}I -mAb A33 compared to ^{131}I -mAb A33 is reduced bone marrow toxicity. With patients who were treated with ^{125}I -mAb A33 at an initial dose of $50\ \text{mCi/m}^2$, no toxicity was observed. Tumor imaging by external ^{125}I scanning with a collimator was achieved in all patients, and positive tumor images were followed for up to 35 days after antibody infusion. This result was surprising since only a small fraction of ^{125}I radiation was previously considered to be suitable for scanning.

35 Most of the ^{125}I radiation acts at very short range, and the positive images suggest significant tumor doses. Of the first nine patients, one showed a response with 35% reduction in carcinoembryonic antigen (CEA) levels at 4 weeks after treatment. It was not possible to evaluate one patient as this patient had a pulmonary lesion resection. Tumor regression resulted from ^{125}I -mAb A33 administration. In addition, patients who did not respond to chemotherapy prior

to being administered ^{125}I -mAb A33 were responsive to chemotherapy (i.e., had tumor regression) after ^{125}I -mAb A33 administration. Therefore, the antibody-radioisotope conjugates of the invention can be used to sensitize patients to chemotherapy.

Example III - Biodistribution of ^{131}I -mAb A33 in Humans with Colon Cancer

The biodistribution of ^{131}I labelled mAb A33 in presurgical patients with suspected hepatic metastases of colorectal cancer was studied. This was done in order to establish the targeting potential of mAb A33 as determined by external scanning techniques and biopsy-based dosimetry, and to show that mAb A33 targeting is antigen-specific when compared to an unrelated isotype-matched negative control mAb, mAb TA99.

Efficient and consistent radiolabelling of mAbs with retention of optimal levels of immunoreactivity is essential for radioimmunotherapy. Preclinical evaluation showed that mAb A33 labelled with up to 50 mCi ^{131}I /mg protein retained 40-70% of the original immunoreactivity as determined by sequential absorption analysis. Fifty-one doses of mAb A33 were labelled in 42 iodination procedures, using a protocol designed to achieve a specific activity of about 15 mCi/mg (actual range 12.5 to 22 mCi/mg). Seventeen patients received samples with 40-70% immunoreactivity. Six patients received some preparations of ^{131}I -mAb A33 at 40-70% immunoreactivity and some with 20-35% immunoreactivity, which is considered to be suboptimal.

Seventeen patients were administered ^{131}I -mAb A33 at five dose levels (0.2-50 mg) and three patients received the unrelated isotype-matched negative control ^{131}I -mAb TA99 (2 mg) together with ^{125}I -mAb A33 (2 mg). Following infusion over a 1-hour period, peak serum levels of ^{131}I -mAb A33 and control ^{131}I -mAb TA99 reached 28% ID/l at all dose levels tested, consistent with free distribution in the intravascular plasma volume (median 2.57 l) as determined from $^{99\text{m}}\text{Tc}$ -HSA serum

studies in the same patients. Clearance of ^{131}I -mAb A33 from the blood showed that the half-life of ^{131}I -mAb A33 was found to decrease more rapidly than the levels of negative control ^{131}I -mAb TA99, most likely due to *in vivo* absorption of ^{131}I -mAb A33 by the antigen-positive tumor tissue.

In the 17 patients treated with ^{131}I -mAb A33, a total of 46 hepatic metastases were documented at surgery, and 45 of these were identified on whole body planar or spot view images of the abdomen. Two primary colon cancers were also visualized by external scans, while regional lymph node metastases, although having similar tumor:normal tissue ratios by biopsy-based dosimetry, could not be distinguished from adjacent sites of uptake in the abdomen.

The vascularity of liver metastases was assessed with $^{99\text{m}}\text{Tc}$ -HSA scans and biopsy-based dosimetry. The $^{99\text{m}}\text{Tc}$ -HSA scans confirmed the hypovascularity characteristic of these lesions, which appeared as photopenic areas compared to the surrounding, uninvolved liver. Similarly, biopsies showed tumor:liver ratios for $^{99\text{m}}\text{Tc}$ ranging from 0.14:1 to 1.19:1, with most lesions having values of less than 0.60. Tests with the negative control mAb, ^{131}I -mAb TA99, showed minimal tumor uptake with faint positive imaging in only one patient, and tumor:liver ratios of 1.1:1 to 5.4:1. In contrast, ^{131}I -mAb A33 produced tumor:liver ratios up to 100:1 as determined from biopsies. Specific accumulation of ^{131}I -mAb A33 in the hepatic lesions but not in surrounding normal liver was also visualized by combined autoradiography (see Figures 1 and 2) and immunohistochemistry with frozen tissue sections.

Variable localization of ^{131}I -mAb A33 to the normal intestine was found. Scans in several patients showed labeling of some portions of the bowel, with clear outlines of the normal colon in some cases. However, in other cases, only minimal normal colon imaging was shown. Tests with sections of normal large bowel removed during surgery showed that a significant amount of radiolabel could be recovered by simply washing the specimens with phosphate-buffered saline. Thus, at least some of the apparent normal bowel uptake of ^{131}I -mAb

A33 may represent free ^{131}I secreted by the stomach or monoclonal antibody if binding affinity was low. The finding that A33-positive normal small and large intestinal mucosa do not accumulate ^{131}I -mAb A33 more consistently may be related to lack of accessibility of the antigen in the normal mucosa.

Example IV - Therapeutic Effect of
 ^{131}I -mAb A33 in Humans with Colon Cancer

A therapy trial of ^{131}I -mAb A33 was conducted in patients with advanced colorectal cancer. Twenty-three patients were entered into the study and received 30 mCi/m² of ^{131}I -mAb A33 as a single infusion or up to three infusions on consecutive days. Two patients (Nos. 14 and 18) were lost to follow-up (insufficient number of blood samples at nadir time) and two patients who died of progressive disease less than one month after treatment (Nos. 2 and 22) had incomplete evaluation of toxicity. Upon review of histologic slides, one patient (No. 10) was found to have small bowel cancer rather than colon cancer. For the patients included in this study, no frozen tissues for A33 immunohistochemistry were obtained, thus precluding antigen typing as a criterion for inclusion in the study. The available mAbs against A33 do not allow antigen detection in routinely fixed, paraffin-embedded material. One to 3 weeks after treatment, 20 of the 23 treated patients showed positive ^{131}I -mAb A33 scans for tumor sites independently identified by computerized tomography (CT) or chest X-rays. Two patients with surgically documented intra-abdominal disease and rising CEA levels were not visualized by either CT or ^{131}I -mAb A33 scan, and one patient (No. 10) with small bowel cancer had a negative ^{131}I -mAb A33 scan.

All patients included in this study developed human anti-mouse antibody ("HAMA") responses after one treatment cycle, with titers of 1:10,000 to 1:100,000 for both IgM and IgG by 4 weeks post-treatment. The major toxicities observed were thrombocytopenia and neutropenia, and their severity was generally dose-dependent. However, for each dose level tested (≥ 45 mCi/m²), patients with prior extensive chemotherapy

showed the most pronounced and/or prolonged signs of bone marrow suppression, even if the platelet counts had returned to the normal range at the time of ^{131}I -mAb A33 treatment. This patient variation in hematologic toxicity was not due to differences in ^{131}I retention as measured by total body isotope clearance rates. Instead, it appeared to reflect an intrinsic sensitivity to radiation damage. Of the five patients treated with 75-84 mCi/m², one patient (No. 11) developed grade 3 thrombocytopenia (nadir 33,000/ μl for 2 days), and three patients (Nos. 13, 14 and 15) displayed no toxicity effects. Patients Nos. 11 and 12 had prior treatment with BCNU, streptozotocin, vincristine, 5-fluorouracil (5-FU), and leucovorin, whereas patients Nos. 13, 14 and 15 received only 5-FU and leucovorin.

Of the six fully evaluable patients who received 86-94 mCi/m² ^{131}I -mAb A33 (Nos. 16, 17, 19, 20, 21 and 23), two (Nos. 21 and 23) showed grade 4 toxicity, one showed grade 1 toxicity, and three showed no toxicity. Of the two patients with grade 4 toxicity, patient No. 21, who had extensive prior chemotherapy (5-FU, interferon- α , BCNU, streptozotocin, and vincristine in separate protocols over a 30-month period), showed rapid and prolonged suppression of platelet and neutrophil counts after ^{131}I -mAb A33 treatment. Neutropenia was seen in two patients (Nos. 21 and 23) with thrombocytopenia. Two additional patients (Nos. 18 and 22) who had only a single blood test during their nadir period, and were therefore not fully evaluable, had at least grade 2 and 3 neutrophil toxicity. Lymphocytopenia generally accompanied decreases in total white cell counts, but was not analyzed with regard to distinct lymphocyte subsets.

Treatment responses were evaluated in 18 patients who had disease which had been measured by either chest X-ray or CT scan, and three patients had disease measurable by serum CEA levels, but failed to show measurable disease by radiographic studies. Two patients were not evaluable due to loss to follow-up or surgical resection of the tumor. Evidence for mixed responses was obtained in five patients (Nos. 3, 5, 7,

9 and 23). Patient No. 3 presented liver metastases and cytology-proven, malignant ascites, which had been stabilized with diuretics (lasix, aldactone) before treatment. After ¹³¹I-mAb A33 infusion, extensive accumulation of radioisotope in the ascitic fluid was detected. One month after treatment, the patient's weight had decreased by 8 kg, the ascites had disappeared, and serum CEA levels had dropped from 3,750-4,200 ng/ml (before treatment) to 2,000 ng/ml. Subsequently, the diuretics were discontinued and serum CEA levels stabilized at 1,750 ng/ml for almost 4 months. The liver metastases remained unchanged by radiographic criteria. Retreatment with ¹³¹I-mAb A33 was attempted after 10 weeks; however, due to HAMA response, isotope accumulation was seen only in liver and spleen, and not in tumor sites.

Patient No. 5 had extensive pleural disease in the right chest, a measurable lesion in the left chest, and liver metastases. Three weeks after ¹³¹I-mAb A33 treatment, the left chest lesion appeared less well-defined, and 2 months later had disappeared (see Figure 3). The other sites of measurable disease remained stable for 5 months. Patient No. 9 had extensive abdominal disease including liver metastases, a large pelvic mass, and large left supraclavicular and neck lymph nodes. Four weeks after ¹³¹I-mAb A33 treatment, the lymph nodes were no longer palpable, although a month later her pelvic mass showed an increase in size. In patients Nos. 7 and 23, serum CEA levels decreased by 22% and 30%, respectively, for more than 16 weeks; however, more detailed assessment of disease progression was not possible since neither patient had measurable disease by radiologic examination. A comparison of the patients with and without evidence of mixed responses has not revealed any consistent differences. In particular, no obvious correlation has emerged between the level of radioisotope uptake into tumor sites (determined from scans) and measurable responses.

Preparation of Antibody Conjugates

Monoclonal antibody A33 can be conjugated to anti-cancer drugs and administered to treat cancer. For example, mAb A33 can be conjugated with QFA, which is an antifolate, or with calicheamicin, which is an anti-tumor antibiotic that cleaves double-stranded DNA of tumor cells. Both QFA and calicheamicin have intracellular sites of action and do not readily cross the plasma membrane. Therefore, they have weak cytotoxic effects when added to cell cultures. Cellular uptake of these agents through mAb A33-mediated internalization greatly enhances their cytotoxic effects *in vitro*. *In vivo* xenograft studies show that tumor inhibition with limited normal tissue damage can be obtained with both mAb A33-QFA and mAb A33-calicheamicin conjugates. Other anti-cancer drugs can also be conjugated to the antibodies of the invention and used to treat cancer, including BCNU, streptozocin, vincristine and 5-fluorouracil.

Monoclonal antibody A33 can also be used to target genetic material to both colon cancer cells and normal colonocytes since it is internalized rapidly into target cells. The A33 antigen directs the antibody to macropinosomes, which are compartments inside the cell. Transfected DNA also traverses the macropinosomal compartment, surviving hydrolysis and ultimately incorporating into chromosomal DNA. With mAb A33-drug conjugates, the A33 antibody directs reagents to the nuclear DNA. Similarly, genetic material can be directed inside the cell utilizing mAb A33.

Example V - A33-Oligonucleotide Conjugate

A 26-mer oligonucleotide containing a free, reactive thiol group at the 3' end was conjugated to a free amine of mAb A33 using the heterobifunctional reagent SPDP. The 5' end of the oligonucleotide was enzymatically labelled with ³²P as tracer for DNA. SDS PAGE and reduced gels showed a pattern consistent with a product containing a range of oligonucleotide/mAb ratios. Purification was accomplished

with protein A chromatography. The final product was sterile-filtered and tested for cell-binding activity and protein and bound oligonucleotide concentrations.

5 The final purified product was tested on an A33 antigen-expressing colon cancer cell line SW1222. Based on the ability of low pH to reverse mAb A33 binding to its antigen, a protocol was developed to quantify the percentage of internalization. This was used to analyze and compare binding and internalization of mAb A33-³²P-DNA with that of ¹²⁵I-mAb
10 A33. An excess of unlabelled and unmodified mAb A33 was used to block the specific binding of radiolabelled mAb A33 conjugates to cells. This demonstrated that all cell binding was specific and mediated through the interaction of mAb A33 and its antigen. The assay was done in 96-well plates with
15 100,000 cells per well in log phase growth. At 24 hours, 7.2% of ³²P added to the wells was bound to the cells. By 48 hours, over 60% of ³²P was internalized. Anti-tumor effects were observed in nude mice carrying transplanted SW1222 cells when anti-actin DNA was used. These conjugates also contained ³²P
20 as a tracer, and it is possible that ³²P contributed somewhat to the anti-tumor effects. However, since the amount of radioactivity was low (10 μ Ci/mouse), minimal effects would be expected from the isotype alone.

25 The antibody conjugates of the invention can be administered via various routes known by those skilled in this field. By way of example, they can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously or orally (e.g., attached to virus). The antibody conjugates of the invention can also be administered
30 in combination with a pharmaceutically acceptable carrier in composition form. The antibody conjugates of the invention can be administered at a dose range of about 0.1 mg to 2000 mg. However, dose ranges will vary among patients to be treated, depending on the condition of the patient to be
35 treated, the medical history of the patient, the means of administration and several other factors.

Example VI - Characterization of the A33 System

In order to characterize the A33 system, several A33-positive colon cancer cell lines were used to determine the number of mAb A33 binding sites per cell. It was found that there are greater than 800,000 sites per cell. The avidity of mAb A33 binding was found to be high. The same cell lines were used to show that A33 is a rapidly internalizing antigen, with up to 90% internalization over a 24-hour period in humans. Studies of the binding kinetics of ¹²⁵I-mAb A33 to cultured colon cancer cells, acid wash experiments, and SDS-gel analysis of cell extracts and culture supernatants showed that 80-85% of the internalized mAb A33 cycles through intracellular compartments and, within about 1-2 days, is released from the cells in intact, immunoreactive form. Only a small fraction of internalized ¹²⁵I-mAb A33 appears to be degraded after internalization, and this degradation was partially blocked by inhibitors of lysosomal enzymes.

The mechanism of mAb A33 internalization involves macropinosomes. *In vitro* studies with mAb A33 conjugated with ¹²⁵I, calicheamicin and QFA showed strong cytotoxic activity against A33-positive colon cancer cells but not against A33-negative tumor cell lines. Since all three conjugates act after intracellular uptake, it was determined that the internalizing capacity of A33 plays an essential role in the observed cytotoxic effects.

Example VII - Generation of Other Murine Antibodies

The inventors have generated two additional mouse mAbs, designated mAb 100-210, which is of the IgG2b isotype, and mAb 100-310, which is of the IgG2a isotype. mAb 100-310 is described in U.S. Patent Application Serial No. 08/273,277, the entirety of which is incorporated herein by reference. Hybridomas secreting monoclonal antibodies 100-210 and 100-310 were deposited under the Budapest Treaty on November 30, 1994 and April 28, 1992, respectively, with the American Type Culture Collection, Rockville, Maryland, and catalogued as ATCC Nos. HB 11764 and HB 11028, respectively. These

monoclonal antibodies show the same patterns of reactivity as mAb A33. Monoclonal antibody 100-210 blocks mAb A33 binding in competitive radiobinding assays. This suggests that it binds to the same or a spatially related epitope on A33 antigen as mAb A33, whereas mAb 100-310 and mAb A33 do not cross-block each other, suggesting that they react with two distinct epitopes on the same cell surface protein on A33 antigen.

Example VIII - Generation of Humanized A33 Antibodies

The use of high- or low-affinity mAbs or mAbs with faster or more delayed serum clearance for immunotherapy in cancer patients will vary for different antigenic targets, tumor types, intrinsic pharmacokinetic properties of the mAbs, intended effector mechanisms, and other nonimmunologic parameters. Since mAb A33 shows excellent targeting properties *in vivo*, it provides an effective test system for evaluating these parameters. Humanized and modified monoclonal antibodies have been produced from mAb A33.

In order to clone and express the genes for mAb for A33, mAb A33 hybridoma cells were cultured in RPMI 1640 medium supplemented with 10% fetal calf serum and 1 mM glutamine. Total RNA was prepared from 10⁹ hybridoma cells using guanadinium isothiocyanate and poly(A+) mRNA isolated by oligo (dT) affinity chromatography. First strand cDNA was synthesized from 10 mg mRNA. DNA sequences encoding A33 variable domains, including signal sequences for secretion, were amplified from the cDNA using the PCR procedure described by Jones et al., Bio/Technology, 9:88-89 (1991), but with primers designed to allow facile cloning of the PCR products into vectors for expression in mammalian cells. These vectors were derived initially from pEE6 (Stevens et al., Nucl. Acids Res., 17:7110 (1989)).

Figure 4 shows final NSO expression vectors. PCR amplified fragments for the light chain variable domain were cloned between the Bst1 and Sph1 sites of pMRR010, a pEE6

derivative constructed to allow expression of such sequences as kappa chimeric light chains. PCR amplified fragments for the A33 heavy chain variable domain were cloned between the HindIII and ApaI sites of pMRR011, a pEE6 derivative constructed to allow expression of such sequences as gamma 1 chimeric heavy chains. The cloned variable region genes were sequenced by the double strand dideoxy chain terminating method using T7 DNA polymerase.

The humanized variable domains were assembled by the procedure disclosed by Daugherty et al., Nucl. Acids Res., 19:2471-2476 (1991) using primers which allowed facile cloning into pMRR010 and pMRR011 for the light and heavy chains, respectively. These humanized variable region genes were sequenced by the same procedures used for the murine variable region genes. An expression vector for the hA33 Fab'Δcys heavy chain with a single hinge thiol was constructed by replacing the gamma 1 constant domains with the appropriate segment from the CB72.3 Fab'Δcys gene described by King et al, Cancer Res., 54:6176-6185 (1994).

Transient co-expression of murine IgG heavy and light chains, or humanized IgG and Fab' heavy and light chains was achieved by co-transfection of the separate expression vectors into CHO-L761 hour cells as described by Cockett et al., Bio/Technology, 8:662-667 (1990). For stable cell line development the heavy and light chain expression, units were combined in a single plasmid. This was accomplished by replacing the NotI-BamHI stuffer fragment in the light chain expression plasmids with the NotI-BamHI fragments carrying the hCMV promoter/enhancer and heavy chain genes from the heavy chain expression plasmids. The final expression plasmids were termed pAL71 and pAL72 for hA33 IgG1 and Fab'Δcys, respectively (Figure 4).

Stable NSO cell lines for the production of hIgG1 and hFab' were then established by transfecting the plasmids according to the procedure taught by Bebbington et al., Bio/Technology, 10:169-175 (1992). After transfection, the cells were plated at 2×10^5 cells per 96-well plate in

Dulbecco's Modified Eagles Medium containing 10% dialysed fetal calf serum and 2 mM glutamine. After 24 hours cells were selected by the addition of methionine sulfoximine to the medium at a final concentration of 7 μ M. After 21 days of culture, resistant colonies were picked and expanded for analysis of productivity. The highest producers were selected for production of recombinant antibody in roller bottle culture.

The results of these binding assays showed that chimeric antibody bound to cells expressing the antigen as well as murine A33, confirming that the cloned genes correspond to those of A33. Figure 5 shows the amino acid sequences of the heavy and light chains deduced from the DNA sequence of the cloned variable domain genes. N-terminal protein sequencing of the first 11 amino acids gave results completely consistent with these deduced amino acid sequences for both heavy and light chains, confirming that the appropriate genes had been cloned.

Figure 5 shows the amino acid sequences of the light [A] and heavy [B] chain variable domains of the murine and humanized A33 antibodies. The sequence of the mouse antibody as deduced from cDNA (mA33) is shown aligned with the humanized antibody sequence (hA33). The humanized framework sequence is derived from the human antibody LAY (Kabat et al., Sequences of Proteins of Immunological Interest, 4th Ed. (1987)). The three complementarity determining regions in each chain are underlined. Residues in the LAY framework that have been replaced with mouse A33 sequence are double underlined.

The V_H of A33 shows closest homology (70%) to the consensus sequence of human subgroup V_H III, while the V_L shows greatest homology to the consensus sequence of human subgroups V_L I and V_L I (62%). It was from these subgroups that LAY, which has a V_H III heavy chain and V_L I light chain, was chosen as the human framework. In Figure 5, the light chain residues 1-23, 35-45, 47-49, 57-86, 88 and 98-108 inclusive were derived from the LAY sequence, and the residues 24-34, 46, 50-56, 87 and

89-97 correspond to the Complementarity Determining Regions (CDRs). Residues 46 and 87 are predicted to be at the interface of the light and heavy variable regions. Residue 46 is usually a leucine in human antibody sequences, and residue 87 is usually a phenylalanine or tyrosine.

For the heavy chain, residues 2-26, 36-49, 66-71, 74-82a, 82c-85, 87-93 and 103 to 113 inclusive were derived from the LAY sequence while residues 1, 27-35, 50-65, 72, 73, 82b, 86 and 94-102 inclusive were derived from the murine sequence. Residues 31-35, 50-65 and 95-102 in the heavy chain correspond to the CDRs. The murine derived amino acids in the framework regions were included for several reasons: residue 1 is usually solvent accessible and in the vicinity of the CDR region (residues 27-20); LAY has a residue, alanine, not normally found at this position in human or murine V_H sequences and therefore the murine residue was used. Also, at positions 72 and 73, the murine residue was used because of the predicted proximity to CDR2 and, in the case of residue 72, to remove the possibility of introducing an N-linked glycosylation site into the variable domain by the use of the LAY framework. The murine sequence was also used at the inter-domain residue 94, where A33 has a proline, not normally found at this position. Murine residues were used at positions 82 and 86 because the use of the human amino acids at these positions in a humanized antibody with LAY frameworks has previously been found to be deleterious for the expression of the heavy chain.

hIgG was purified from tissue culture supernatants of NSO cells using protein A-Sepharose affinity chromatography and characterized by SDS-PAGE as described by King et al., Biochem., 281:317-323 (1992). hA33 Fab' Δ cys was purified from cell culture supernatant by chromatography on protein A-Sepharose using the low affinity protein A binding site on Fab' as a basis for purification. A column of protein A-Sepharose was equilibrated with 100 mM boric acid buffer pH 8.0 containing 150 mM sodium chloride. The tissue culture supernatant from NSO cells expressing hA33 Fab' Δ cys was

adjusted to pH 8.0 by the addition of 1 M tris and applied to the column. After washing with the equilibration buffer, the Fab' was eluted with 0.1 M citric acid, collecting fractions directly into sufficient 1 M tris to immediately adjust the pH of the fraction to between 6 and 7.

After dialysis and ultrafiltration, hA33 Fab'Δcys was cross-linked to DFM™ and TFM™ using the maleimide-based homobifunctional and homotrifunctional linkers CT52 and CT998 respectively, as described for cB72.3 Fab'Δcys by King et al., (1994) supra. These cross-linkers contain the 12N4 macrocycle for incorporation of ⁹⁰Y. The DFM™ and TFM™ produced were purified by gel filtration using a Sephacryl S-200HR column. Radiolabelling with ⁹⁰Y was performed as described by King et al., (1994) supra.

Small amounts of the humanized antibody were produced in a transient expression system in CHO cells to establish that it bound SW1222 cells expressing the antigen. Stable NSO cell lines were then isolated to produce larger quantities of purified material, for both hA33 IgG and Fab'Δcys. The best lines produced approximately 700 mg/ml of hA33 IgG1 and 500 mg/ml of Fab'Δcys in suspension culture.

Figure 6 shows SDS polyacrylamide gel of humanized A33 IgG and TFM™ under (A) non-reducing and (B) reducing conditions. Lane 1, hA33 IgG; lane 2, hA33 Fab'Δcys; lane 3 hA33 Fab'Δcys cross-linking mix; lane 4 purified hA33 TFM™.

Figure 6 demonstrates that the purified hIgG was homogeneous and fully assembled. HPLC analysis demonstrated that it was free of aggregates. As expected, the hFab'Δcys was recovered largely in the form of monovalent Fab' with little in the form of F(ab')₂, consistent with results for other recombinant Fab' fragments. SDS-PAGE analysis of the purified TFM™ showed a single species of approximately 150 kD under non-reducing conditions. Under reducing conditions, two species of approximately 75 kD and 25 kD were observed, corresponding to the sizes expected for three cross-linked Fd' chains and light chains, respectively.

Cross-linking of Fab'Δcys to TFM™ was achieved with a

yield of 60-65%. Figure 7 shows HPLC profiles at 280 nm of (a) hA33 Fab'; (b) hA33 Fab' after cross-linking to TFM[™] and (c) purified hA33 TFM[™]. HPLC gel filtration was carried out on a DuPont Zorbax GF-250 column run at 1 ml/min in 0.2 M sodium phosphate buffer pH 7.0. As shown in (a) hA33 Fab' peak has a retention time of 10.5 minutes with a minor peak of F(ab')₂ at 9.7 minutes and a buffer peak at 12.2 minutes. As shown in (b) after cross-linking, the major peak represents TFM[™] at 9.0 minutes. Minor peaks represent di-Fab at 9.6 minutes, residual monomeric Fab' at 10.5 minutes, a small amount of aggregate at 8.2 minutes and the same buffer peak at 12.2 minutes. As shown in (c), after purification, the TFM[™] peak is still seen at 9.0 minutes, the only other visible peak being the buffer peak at 12.2 minutes.

Antigen binding assays were then performed using cells of two human colorectal tumor cell lines which express the A33 antigen, COLO 205 and SW1222. These cells were cultured in DMEM medium containing 2 mM glutamine and 10% fetal calf serum. Assembled antibody in culture supernatants and in purified preparations was quantitated in an ELISA. Direct binding of murine and humanized antibodies in SW1222 cells was measured in a FACScan assay. SW1222 cells were trypsinized to remove them from culture flasks, washed with phosphate buffered saline (PBS) and resuspended in PBS containing 10% bovine serum and 0.1% sodium azide. Humanized A33 antibody was serially diluted in PBS containing 10% FCS and 0.1% sodium azide and added to 2×10^5 cells. Following a one hour incubation on ice, the cells were washed in PBS and then incubated with a rhodamine anti-(human Fc) conjugate (1:1000 in PBS, 10% FCS and 0.1% sodium azide) for one further hour on ice. After washing in PBS, the amount of rhodamine labelled A33 antibody conjugate bound to the cells was measured in a FACScan analyzer. Direct binding of murine A33 was measured by FACScan analysis after incubation of SW1222 cells with FITC-labelled antibody. Suitable non-specific antibody controls were carried out to demonstrate that A33 binds specifically via antibody-antigen interaction rather than non-

specifically through Fc interactions.

Determination of affinities of murine and humanized antibodies was based on the procedure described by Krause et al., Behring Inst. Mitt., 87:56-67 (1990). Briefly, antibodies were labelled with fluorescein using fluorescein isothiocyanate (FITC) titrated from 1.3 mg/ml, and then incubated with 2.8×10^5 SW1222 cells for two hours on ice in 350 ml PBS containing 5% FCS and 0.1% sodium azide. The amount of fluorescence bound per cell was determined in a FACScan and calibrated using standard beads. The number of molecules of antibody that had bound per cell at each antibody concentration was thus established and used to generate Scatchard plots. Competition assays were performed by FACScan quantitation of bound FITC-labelled murine A33 after incubating SW1222 cells with a standard quantity of the murine antibody together with a dilution series of the humanized variants.

Figure 8 shows Scatchard analysis for the murine antibody and hIgG1 binding to SW1222 cells. These data suggest that both antibody forms have equilibrium dissociation constants (K_D 's) of 1.3 nM and have approximately 300,000 sites per cell. The antigen binding activity of hTFM[™] was compared to those of monovalent hFab' Δ cys and hIgG in competition binding assays in which these species were competed with murine IgG for binding to COLO 205 cells expressing the antigen. The results (Figure 9) demonstrate that the monovalent Fab' fragment binds less well than the bivalent IgG, as expected. The trivalent hTFM[™], on the other hand, showed approximately twofold better binding than hIgG, presumably as a result of increased avidity. This finding is consistent with results for chimeric B72.3, for which TFM[™] also showed two to threefold better binding than IgG.

Figure 8 shows a Scatchard plot of humanized and murine A33 binding to SW1222 cells. Experimental details as in materials and methods. K_D values of 1.28 nM for murine A33 IgG and 1.27 nM for humanized A33 IgG were calculated from linear regression analysis of the data points. Figure 9

represents the competitive binding assay for hA33 Fab' Δ cys, IgG and TFM[™] binding to COLO 205 cells. hA33 IgG (\diamond), TFM[™] (|), DFM[™] (\square), and Fab' Δ cys (Δ) were competed with FITC labelled murine A33 IgG and results plotted as fluorescence units versus nM binding sites.

Antibodies were then labelled with ⁹⁰Y via the macrocyclic ligand tetraazocyclododecane etra-acid, (termed 12N4 or DOTA) coupled to the immunoglobulin via 12N4-maleimide linkers as described by Antoniow et al., "Practical evaluation of yttrium-90 labelled monoclonal antibody A33 and A33 tri-fab (TFM) for radioimmunotherapy of colorectal carcinoma", In Press. Radiolabelling of hA33-12N4 conjugates with ⁹⁰Y and ¹²⁵I, and biodistribution studies in nude mice bearing subcutaneous SW1222 tumor xenografts, were carried out as described by Antoniow et al., supra. Antibodies were labelled with ¹¹¹In via a second macrocyclic ligand, 1,4,7-triazacyclononanetriacetic acid or 9N3 using 9N3-maleimide linkers as described by Turner et al., Br. J. Cancer, 70:35-41 (1994). The immunoconjugate was shown to be fully immunoreactive using the competition based FACs assay both before and after radiolabelling. For biodistribution experiments, radiolabelling was achieved to a specific activity of 2 μ Ci/ μ g with >95% incorporation of ⁹⁰Y.

Biodistribution studies in guinea pigs were carried out after i.v. administration to male outbred Dunkin-Hartley guinea pigs of approximately 250-300 grams. Groups of four guinea pigs were injected with each ⁹⁰Y-labelled component into the ear vein and sacrificed post administration at the time intervals indicated in Figure 10. Blood samples were taken and tissues processed as described for mice by Antoniow et al., supra. Pharmacokinetic studies in cynomolgous monkeys of 5-7 kg (2 per group) were also carried out after i.v. injection of radiolabelled components. Blood samples were taken at 0.5, 1, 2, 4, 6, 8, 24, 48, 72, 96, 120, 144 and 168 hours for counting.

Figure 10 shows the time course study for biodistribution of ⁹⁰Y-labelled humanized A33 in nude mice bearing SW1222 tumor xenografts. Mice were injected i.v. with 19 μ Ci (10 μ g) ⁹⁰Y-

labelled hA33 each. Groups of 4 mice were killed at 3 (■), 24 (□), 48 (□) and 144 hours (□) post administration, and the amount of activity was determined in tumor and normal tissues. Each column represents the mean obtained from 4 mice. Error bars represent the standard deviation of the mean. Tissue uptake is plotted as % injected dose per gram of tissue. A favorable biodistribution was achieved, with high levels of activity localized to the tumor and little or no accumulation in any normal tissue. The biodistribution of the humanized A33 immunoconjugate was not significantly different from that of the murine antibody in the same xenograft system at these time points. For both the murine and humanized antibodies the level of activity localized to the tumor increased with time, even though levels in all other tissues were falling, which led to increasing tumor to normal tissue ratios over time (Table 2).

To assess whether this was a feature of the antibody itself or the radioisotope used, a biodistribution experiment was also carried out with humanized A33 labelled with ^{125}I . In this experiment, the absolute levels of isotope retained by the tumor were slightly lower but the tumor to blood ratios were very similar, suggesting that the increasing localization is a property of the A33 antibody rather than the nature of the isotope/chelator system. The lower absolute levels of ^{125}I -hA33 localized to the tumor are probably the result of dehalogenation of the radioiodinated antibody.

TABLE 2**(a) ⁹⁰Y Humanized A33****Time Post Injection**

		3 h	24 h	48 h	144 h
5	BLOOD	0.32±0.08	2.77±0.38	5.63±1.84	10.1±2.60
	MUSCLE	12.8±4.95	34.3±5.93	48.6±8.35	127.3±38.0
	BONE	5.31±2.27	24.4±2.14	32.4±11.2	68.5±21.4
	LUNG	1.29±0.56	9.81±2.03	13.7±1.02	26.5±8.39
	LIVER	1.50±0.37	9.50±1.28	11.8±4.54	30.7±9.02
10	SPLEEN	1.77±0.92	10.1±1.42	12.8±4.91	20.7±5.47
	KIDNEY	1.33±0.25	7.75±1.25	10.3±2.17	21.3±5.60
	COLON	3.90±1.04	19.2±2.78	25.7±10.5	56.7±14.5

(b) ¹²⁵I Humanized A33**Time Post Injection**

		3 h	24 h	48 h	168 h
	BLOOD	0.36±0.05	2.44±0.09	4.75±0.83	8.92±2.12
	MUSCLE	10.8±3.47	27.3±10.2	42.7±9.67	75.5±19.9
	BONE	4.86±1.20	28.9±8.88	41.5±9.74	74.5±21.5
20	LUNG	1.39±0.26	8.16±0.59	14.2±2.24	25.1±6.78
	LIVER	1.12±0.08	8.01±1.80	17.8±3.61	32.8±9.13
	SPLEEN	1.67±0.31	17.3±2.54	29.9±6.12	45.7±18.5
	KIDNEY	1.23±0.21	9.92±0.78	18.2±3.40	33.2±9.88
	COLON	5.30±1.38	22.8±4.00	33.5±6.34	60.5±15.9

25

30

35

It was consistently observed that humanized TFM™ and all other humanized fragments examined clear aberrantly quickly from the circulation of mice, far more quickly than the equivalent murine fragments. This phenomenon was shown to be specific to mice and did not occur in rats, guinea pigs or monkeys. Table 3 shows a comparison of biodistribution for hA33 IgG, DFM™ and TFM™ in guinea pigs. It demonstrates the more rapid blood clearance of the DFM™ and TFM™, with blood activities falling to 0.01 and 0.02% i.d./g respectively at the 144 hours time point. Blood activity for hIgG was much higher, at 0.4% i.d./g, at this time point. Table 3

demonstrates very clearly that for DFM™, much higher levels of radioactivity are taken up by the kidney than for IgG and TFM™. This high activity for the DFM™ clears much more slowly from the kidney than from the blood. At early time points, kidney levels were a little higher for TFM™ and IgG, but much lower than for DFM™, and the activity cleared much faster from the kidney for TFM™ than for DFM™. These results for A33 are consistent with those for DFM™ and TFM™ of cB72.3, and are consistent with the view that the kidney is the organ of clearance for TFM™.

TABLE 3
(a) ⁹⁰Y Humanized A33

		Time Post Injection			
		24 h	48 h	72 h	144 h
	BLOOD	0.88±0.04	0.75±0.06	0.47±0.12	0.40±0.09
	MUSCLE	0.07±0.01	0.07±0.005	0.04±0.01	0.04±0.01
	BONE	0.15±0.01	0.13±0.02	0.08±0.02	0.07±0.02
	LUNG	0.29±0.02	0.25±0.02	0.16±0.05	0.17±0.05
	LIVER	0.27±0.08	0.24±0.02	0.17±0.04	0.16±0.04
	SPLEEN	0.30±0.06	0.31±0.03	0.26±0.06	0.20±0.02
	KIDNEY	0.27±0.003	0.27±0.02	0.18±0.05	0.20±0.05
	COLON	0.12±0.01	0.12±0.01	0.08±0.02	0.07±0.14

(b) ⁹⁰Y hA33 TFM™

		Time Post Injection			
		24 h	48 h	72 h	144 h
	BLOOD	0.58±0.10	0.22±0.05	0.12±0.01	0.02±0.005
	MUSCLE	0.03±0.01	0.02±0.007	0.01±0.001	0.01±0.001
	BONE	0.16±0.03	0.06±0.02	0.04±0.01	0.01±0.002
	LUNG	0.28±0.04	0.12±0.03	0.08±0.01	0.02±0.003
	LIVER	0.28±0.05	0.14±0.05	0.09±0.01	0.04±0.006
	SPLEEN	0.22±0.03	0.11±0.04	0.07±0.01	0.04±0.003
	KIDNEY	1.08±0.25	0.58±0.22	0.49±0.04	0.23±0.03
	COLON	0.13±0.03	0.5±0.01	0.04±0.004	0.02±0.001

TABLE 3 (con't)(c) ^{90}Y hA33 DFM[™]

		Time Post Injection			
		24 h	48 h	72 h	144 h
5	BLOOD	0.17±0.10	0.08±0.01	0.04±0.05	0.01±0.002
	MUSCLE	0.03±0.002	0.02±0.002	0.02±0.03	0.01±0.002
	BONE	0.10±0.01	0.08±0.01	0.07±0.01	0.06±0.012
	LUNG	0.14±0.01	0.09±0.01	0.06±0.01	0.04±0.01
	LIVER	0.28±0.03	0.22±0.03	0.18±0.03	0.12±0.02
10	SPLEEN	0.26±0.01	0.26±0.06	0.21±0.06	0.16±0.04
	KIDNEY	9.31±1.56	8.60±0.98	5.97±1.41	4.27±0.37
	COLON	0.11±0.02	0.07±0.01	0.07±0.01	0.05±0.01

An exponential, nonlinear least-squares fitting procedure was used to determine the blood clearance parameters for each monkey. Mean values of the blood clearance parameters for IgG, TFM[™] and DFM[™] were used to set up appropriate integrals to calculate the percent of absorbed dose as a function of time post administration. Estimates were made as to the absorbed dose for red marrow which would be delivered by each form of the antibody (IgG, TFM[™] and DFM[™]) in humans when labelled with ^{90}Y . The monkey data showed that at early times, all of the administered activity was in the blood circulation, and this was taken to be the case for humans. Calculations were made based on the assumptions that the pharmacokinetics of ^{111}In and ^{90}Y labelled antibodies are the same as each other and the same in monkeys and humans. It was also assumed that there is no specific uptake or radiolabelled antibodies in the marrow so that the radioactivity in the blood and marrow are the same after a few (<5) hours. To generate numbers representative of humans, the following data for standard man were used: a total blood volume of 5000 ml; marrow spaces; absorbed fractions for ^{90}Y beta-particles; and the thickness of the endosteal layer. Due to the high energy of the ^{90}Y beta-particles, the radiation absorbed dose in the marrow and the endosteal layer are for all practical purposes the same.

Figure 11 represents the pharmacokinetic profiles of ^{111}In hA33 IgG (■), TFM[™] (○) and DFM[™] (▲) in cynomolgus monkeys. Data was corrected for decay and plotted as mean percentage injected dose remaining at each time point. Due to safety considerations, these components were labelled with ^{111}In rather than ^{90}Y , since previous work suggested that ^{90}Y -labelled IgG and fragments show pharmacokinetics and biodistribution very similar to those labelled with ^{111}In . The plasma clearance profiles are shown in Figure 11 with the alpha and beta phase half life values in Table 4.

TABLE 4

Antibody Form	t1/2 α (hours)	t1/2 β (hours)
IgG	15.8	129.0
TFM [™]	12.8	53.7
DFM [™]	10.8	42.0

As expected from pharmacokinetic data in ice and guinea pigs, both TFM[™] and DFM[™] cleared faster from the circulation than IgG. In addition, DFM[™] cleared more quickly than TFM[™]. When plasma clearance was examined without decay correction for the isotope (Table 5) the data was most consistent with monophasic kinetics for IgG and TFM[™] with biphasic kinetics for DFM[™]. Dosimetric calculations based on these data suggest that when labelled with ^{90}Y at equivalent amounts of radioactivity injected, the DFM[™] gives approximately a five-fold lower absorbed dose to the bone marrow than IgG, while TFM[™] gives a two-fold lower absorbed dose (Table 5).

Table 5

Antibody Form	Effective half life (hours)	Absorbed dose to red marrow & endosteum (rads/mCl)
IgG	40.5	12.0
TFM TM	21.9	6.3
DFM TM	4.7 α /23.6 β	2.6

Single chain antibodies (SCA) can also be generated from mAb A33. Single chain antibodies are small in size, which allows for better diffusion into tumor tissues. They also have a short serum half-life, which reduces toxicity, for example, to the bone marrow. Lower tumor uptake may result from rapid SCA clearance from the blood. Single chain antibody SCA-A33 was produced. This antibody retains the binding specificity of mAb A33 and shows about 3-fold lower binding affinity than the intact mAb. SCA-A33, as well as other single chain antibodies, may be used therapeutically for local/regional perfusion via the hepatic artery for liver metastases and for primary lesions via superior mesenteric artery perfusion.

Although the invention herein has been described with reference to particular embodiments, it is to be understood that these embodiments are merely illustrative of various aspects of the invention. Thus, it is to be understood that numerous modifications may be made in the illustrative embodiments and other arrangements may be devised without departing from the spirit and scope of the invention.

WE CLAIM:

1. A method of reducing the effects of colon cancer in a subject comprising administering to said subject a pharmaceutically effective amount of an anti-cancer drug conjugated to an antibody, wherein said antibody is specific for said colon cancer.

2. The method of Claim 1 wherein said antibody is monoclonal antibody.

3. The method of Claim 2 wherein said monoclonal antibody is selected from the group consisting of monoclonal antibodies A33, 100-210 and 100-310.

4. The method of Claim 1 wherein said antibody is selected from the group consisting of a humanized antibody, a chimeric antibody, a trimeric antibody, a heteromeric antibody and a single chain antibody.

5. The method of Claim 1 wherein said anti-cancer drug is selected from the group consisting of calicheamicin, QFA, BCNU, streptozoicin, vincristine and 5-fluorouracil.

6. A method of reducing the effects of colon cancer in a subject comprising administering to said subject a pharmaceutically effective amount of a conjugate of an antibody specific for said colon cancer and a peptide, wherein said peptide specifically inhibits DNA activity of said colon cancer.

7. The method of Claim 6 wherein said antibody is a monoclonal antibody.

8. The method of Claim 7 wherein said monoclonal antibody is selected from the group consisting of monoclonal antibodies A33, 100-210 and 100-310.

9. The method of Claim 6 wherein said antibody is selected from the group consisting of a humanized antibody, a chimeric antibody, a trimeric antibody, a heteromeric antibody and a single chain antibody.

10. A method of reducing the effects of colon cancer in a subject comprising administering to said subject a pharmaceutically effective amount of a conjugate of an antibody specific for said colon cancer and a radioisotope.

11. The method of Claim 10 wherein said antibody is a monoclonal antibody.

12. The method of Claim 11 wherein said monoclonal antibody is selected from the group consisting of monoclonal antibodies A33, 100-210 and 100-310.

13. The method of Claim 10 wherein said antibody is selected from the group consisting of a humanized antibody, a chimeric antibody, a trimeric antibody, a heteromeric antibody and a single chain antibody.

14. The method of Claim 10 wherein said radioisotope is selected from the group consisting of ^{125}I , ^{131}I , ^{99}Tc , ^{90}Y and ^{111}In .

15. A method of delivering genetic material to DNA of colon cancer cells comprising contacting said cells with genetic material conjugated to an antibody which is internalized into said cells.

16. The method of Claim 15 wherein said antibody is selected from the group consisting of monoclonal antibody A33, monoclonal antibody 100-210 and monoclonal antibody 100-310.

17. A method of delivering an anti-cancer agent to DNA of colon cancer cells comprising contacting said cells with an anti-cancer agent conjugated to an antibody which is internalized into said cells.

18. The method of Claim 17 wherein said antibody is selected from the group consisting of monoclonal antibody A33, monoclonal antibody 100-210 and monoclonal antibody 100-310.

19. The method of Claim 17 wherein said anti-cancer agent is selected from the group consisting of calicheamicin, QFA, BCNU, streptozocin, vincristine and 5-fluorouracil.

20. A peptide-antibody conjugate which is internalized into colon cancer cells when put into contact with said cells.

21. An anti-cancer agent-antibody conjugate which is integrated into colon cancer cells when put into contact with said cells.

22. Hybridoma secreting monoclonal antibody 100-210 deposited on November 30, 1994 with the American Type Culture

Collection, Rockville, Maryland and catalogued as ATCC No. HB 11764.

23. Monoclonal antibody secreted by hybridoma HB 11764.

24. Humanized antibody produced utilizing monoclonal antibody specific for colon cancer cells.

25. Chimeric antibody produced utilizing monoclonal antibody specific for colon cancer cells.

26. Trimeric antibody produced utilizing monoclonal antibody specific for colon cancer cells.

27. Single chain antibody produced utilizing monoclonal antibody specific for colon cancer cells.

FIG. 1





FIG. 2A





FIG. 2B



FIG. 3



3/12

FIG. 4A**FIG. 4B**

4/12


FIG. 5A

mA33	DIVMTQSQKFMSTSVGDRVSITCKASONV
hA33	DIQMTQSPSSLVSVGDRVTITCKASONV
mA33	TLIYLASNRRHTGVPDRFTGSGGTDFTLTISNVQSEDLADYFCLOHWSYP
hA33	TLIYLASNRRHTGVPDRFTGSGGTDFTFTISSLPEDIAFYFCQOHWSTP
mA33	LTFGSGTKLEVKR
hA33	LTFGQGTKEVVR

FIG. 5B


mA33	EVKLVEGGG LVKPGGSLKL SCAASGF
hA33	EVQLLEGGG LVQPGGSLRL SCAASGF
mA33	ISSGGSYTYLLDSVKGRFTISRDSARNTLYLQMSLRSED TALYYCAPTT
hA33	ISSGGSYTYLLDSVKGRFTISRDSARNTLYLQMNLSLQAEI SAIYYCAPTT
mA33	VVPFAYWGQGT
hA33	VVPFAYWGQGT

FIG. 6A



1 2 3 4


FIG. 6B



1 2 3 4


6/12

FIG. 7A




7/12

FIG. 7B




8/12

FIG. 7C




9/12

FIG. 8




10/12

FIG. 9




11/12

FIG. 10



12/12

FIG. 11



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/07321

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 39/395, 48/00; C07K 16/30, 16/46; C12N 5/20

US CL : 424/178.1, 183.1; 530/387.3, 391.3, 391.7; 435/240.27; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/178.1, 183.1; 530/387.3, 391.3, 391.7; 435/240.27; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APL, CAS ONLINE, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCIENCE, Volume 261, issued 09 JULY 1993, Trail et al., "Cure of Xenografted Human Carcinomas by BR96-Doxorubicin Immunoconjugates", pages 212-215, see entire document.	1, 2, 4, 17, 21, 24, 25
X --- Y	PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, Volume 34, issued March 1993, Welt et al., "Localization of ¹¹¹ In-Monoclonal Antibody A33 to Human Colon Cancer Transplants in nu/nu Mice", page 476, abstract no. 2843, see entire abstract.	1-3, 10-12, 14, 17, 18, 21 ----- 4-9, 13, 15, 16, 19, 20, 22-27
X --- Y	JOURNAL OF CLINICAL ONCOLOGY, Volume 12, Number 8, issued August 1994, Welt et al., "Phase I/II Study of Iodine 131-Labeled Monoclonal Antibody A33 in Patients with Advanced Colon Carcinoma", pages 1561-1571, see entire document.	1-3, 10-12, 14, 17, 18, 21 ----- 4-9, 13, 15, 16, 19, 20, 22-27

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

05 JULY 1996

Date of mailing of the international search report

22 JUL 1996

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PAULA HUTZELL

Telephone No. (703) 303-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/07321

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ANTI-CANCER DRUGS, Volume 5, issued 1994, Sondak et al., "Didemnin B in Metastatic Malignant Melanoma: a Phase II Trial of the Southwest Oncology Group", pages 147-150, see entire document.	6-9, 17, 18, 20, 21
Y	COMPARATIVE BIOCHEMISTRY AND PHYSIOLOGY, Volume 105C, Number 20, issued 1993, Yamazaki, "Antitumor and Antimicrobial Glycoproteins", pages 141-146, see entire document.	6-9, 17, 18, 20, 21
Y	US, A, 5,391,723 (PRIEST) 21 February 1995, see entire document.	15
Y	US, A, 4,861,579 (MEYER, JR. ET AL.) 29 August 1989, see entire document.	5, 17, 19
Y	WO, A, 90/07861 (PROTEIN DESIGN LABS, INC.) 26 July 1990, see entire document.	4, 9, 13, 24, 25
Y	WO, A, 94/04691 (THE DOW CHEMICAL COMPANY) 03 March 1994, see entire document.	4, 9, 13, 26, 27

THIS PAGE BLANK (USPTO)

This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**

THIS PAGE BLANK (USPTO)